arXiv:0704.2262v1 [math.NT] 18 Apr 2007

# Irreducible representations and Artin $L$-functions of quasi-cyclotomic fields

Sunghan Bae$^{(1)}$, Yong Hu$^{(2)}$, Linsheng Yin$^{(2)}$
(1) Department of Mathematics, KAIST, Daejeon, Korea
(2) Dept. of Math. Sciences, Tsinghua Univ., Beijing 100084, China

February 1, 2008

**Abstract**

We determine all irreducible representations of primary quasi-cyclotomic fields in this paper. The methods can be applied to determine the irreducible representations of any quasi-cyclotomic field. We also compute the Artin $L$-functions for a class of quasi-cyclotomic fields.

## 1 Introduction

A quadratic extension of a cyclotomic field, which is non-abelian Galois over the rational number field $\mathbb{Q}$, is called a quasi-cyclotomic field. All quasi-cyclotomic fields are described explicitly in [8] followed the works in [1] and [3]. They are generated by a canonical $\mathbb{Z}/2\mathbb{Z}$-basis. The minimal quasi-cyclotomic field containing the quadratic roots of one element of the basis is called a primary quasi-cyclotomic field. L.Yin and C.Zhang [7] have studied the arithmetic of any quasi-cyclotomic field. In this paper we determine all irreducible representations of primary quasi-cyclotomic fields. The methods apply to determine the irreducible representations of an arbitrary quasi-cyclotomic field. We also compute the Artin $L$-functions for a class of quasi-cyclotomic fields.

First we recall the constructions of primary quasi-cyclotomic fields. Let $S$ be the set consisting of $-1$ and all prime numbers. For $p \in S$, we put $\bar{p} = 4, 8, p$ and set $p^* = -1, 2, (-1)^{\frac{p-1}{2}}p$ if $p = -1, 2$ and an odd prime number, respectively. Let $K = \mathbb{Q}(\zeta_{\bar{p}q})$ be the cyclotomic field of conductor $\bar{p}q$. For a class $[a] \in \mathbb{Q}/\mathbb{Z}$, we set $\sin[a] = 2\sin a\pi$ for $0 < a < 1$ and

$\sin[0] = 1$. For prime numbers $p < q$, we define

$$v_{pq} = \prod_{i=0}^{\frac{p-1}{2}} \prod_{j=0}^{\frac{q-1}{2}} \frac{\sin[\frac{iq+j}{pq}]}{\sin[\frac{jp+i}{pq}]} \qquad (p > 2)$$

and

$$v_{2q} = \frac{\sin[\frac{1}{4}]}{\sin[\frac{1}{4q}]} \prod_{j=0}^{\frac{q-1}{2}} \frac{\sin[\frac{j}{2q}] \cdot \sin[\frac{2j-1}{4q}]}{\sin[\frac{4j+i}{4q}] \cdot \sin[\frac{j}{q}] \cdot \sin[\frac{2j-1}{2q}]}.$$

For $p < q \in S$, we put

$$u_{pq} := \begin{cases} \sqrt{q^*} & \text{if } p = -1 \\ v_{pq} & \text{if } p = 2 \text{ or } p \equiv q \equiv 1 \bmod 4 \\ \sqrt{p} \cdot v_{pq} & \text{if } p \equiv 1,\ q \equiv 3 \bmod 4 \\ \sqrt{q} \cdot v_{pq} & \text{if } p \equiv 3,\ q \equiv 1 \bmod 4 \\ \sqrt{pq} \cdot v_{pq} & \text{if } p \equiv q \equiv 3 \bmod 4. \end{cases}$$

Let $\tilde{K} = K(\sqrt{u_{pq}})$. Then $\tilde{K}$ is the minimal one in all quasi-cyclotomic fields which contain $\sqrt{u_{pq}}$. We call these fields $\tilde{K}$ primary quasi-cyclotomic fields. Let $G = \mathrm{Gal}(K/\mathbb{Q})$ and $\tilde{G} = \mathrm{Gal}(\tilde{K}/\mathbb{Q})$. We always denote by $\epsilon$ the unique non-trivial element of $\mathrm{Gal}(\tilde{K}/K)$. If $(p, q) = (-1, 2)$, then the group $G$ is generated by two elements $\sigma_{-1}$ and $\sigma_2$, where $\sigma_{-1}(\zeta_8) = \zeta_8^{-1}$ and $\sigma_2(\zeta_8) = \zeta_8^5$. If $p > 2$, then $G$ is generated by two elements $\sigma_p$ and $\sigma_q$, where $\sigma_p(\zeta_p) = \zeta_p^a, \sigma_p(\zeta_q) = \zeta_q$ and $\sigma_q(\zeta_p) = \zeta_p, \sigma_q(\zeta_q) = \zeta_q^b$, with $a, b$ being generators of $(\mathbb{Z}/p\mathbb{Z})^*$ and $(\mathbb{Z}/q\mathbb{Z})^*$ respectively. If $p = 2$, then $G$ is generated by three elements $\sigma_{-1}, \sigma_2$ and $\sigma_q$, where $\sigma_{-1}, \sigma_2$ act on $\zeta_8$ as above and on $\zeta_q$ trivially, and $\sigma_q$ acts on $\zeta_q$ as above and on $\zeta_8$ trivially.

Next we describe the group $\tilde{G}$ by generators and relations. An element $\sigma \in G$ has two liftings in $\tilde{G}$. By [Sect.3, 7] the action of the two liftings on $\sqrt{u_{pq}}$ has the form $\pm\alpha\sqrt{u_{pq}}$ or $\pm\alpha\sqrt{u_{pq}}/\sqrt{-1}$ with $\alpha > 0$. We fix the lifting $\tilde{\sigma}$ of $\sigma$ to be the one with the positive sign. Then the other lifting of $\sigma$ is $\tilde{\sigma}\epsilon$. The group $\tilde{G}$ is generated by $\epsilon, \tilde{\sigma}_p$ and $\tilde{\sigma}_q$ (and $\tilde{\sigma}_{-1}$ if $p = 2$). Clearly $\epsilon$ commutes with the other generators. In addition, we have $\tilde{\sigma}_p\tilde{\sigma}_q = \tilde{\sigma}_q\tilde{\sigma}_p\epsilon$ (and $\tilde{\sigma}_{-1}$ commutes with $\tilde{\sigma}_2$ and $\tilde{\sigma}_q$ if $p = 2$). For an element $g$ of a group, we denote by $|g|$ the order of $g$ in the group. Let $\log_{-1} : \{\pm 1\} \to \mathbb{Z}/2\mathbb{Z}$ be the unique isomorphism. For an odd prime number $p$ and an integer $a$ with $p \nmid a$, let $(\frac{a}{p})$ be the quadratic residue symbol. We also define $(\frac{a}{2}) = (\frac{a}{-1}) = 1$ for any $a$. Then we have, see [7, Th.3],

$$|\tilde{\sigma}_p| = (1 + \log_{-1}(\frac{q^*}{p}))|\sigma_p| \quad \text{and} \quad |\tilde{\sigma}_q| = (1 + \log_{-1}(\frac{p^*}{q}))|\sigma_q|,$$

with the exception that $\tilde{\sigma}_2 = 2|\sigma_2|$ when $(p, q) = (-1, 2)$. If $p = 2$, we have furthermore $|\tilde{\sigma}_{-1}| = |\sigma_{-1}|$. Thus we have determined the group $\tilde{G}$ by generators and relations.

# 2 Abelian subgroup of index 2

In this section we construct a subgroup of $\tilde{G}$ of index 2 and determine the structure of the subgroup. We consider the following three cases separately:

Case A: $|\widetilde{\sigma}_p| = |\sigma_p|$ and $|\widetilde{\sigma}_q| = |\sigma_q|$;
Case B: $|\widetilde{\sigma}_p| = 2|\sigma_p|$, $|\widetilde{\sigma}_q| = |\sigma_q|$ or $|\widetilde{\sigma}_p| = |\sigma_p|$, $|\widetilde{\sigma}_q| = 2|\sigma_q|$;
Case C: $|\widetilde{\sigma}_p| = 2|\sigma_p|$ and $|\widetilde{\sigma}_q| = 2|\sigma_q|$.

All the three cases may happen. In fact, the case (A) happens if and only if $(\frac{p^*}{q}) = (\frac{q^*}{p}) = 1$; the case (B) happens if and only if $(\frac{p^*}{q}) \neq (\frac{q^*}{p})$ or $(p,q) = (-1,2)$; and the case (C) happens if and only if $(\frac{p^*}{q}) = (\frac{q^*}{p}) = -1$.

In the case A, we define the subgroup $N$ of $\widetilde{G}$ to be

$$N = \begin{cases} < \widetilde{\sigma}_{-1}, \widetilde{\sigma}_2, \widetilde{\sigma}_q^2, \varepsilon > & \text{if} \quad p = 2 \\ < \widetilde{\sigma}_p, \widetilde{\sigma}_q^2, \varepsilon > & \text{if} \quad p \neq 2 \end{cases} \tag{A2.1}$$

It is easy to see that the subgroup $N$ is abelian of index 2 and is a direct sum of the cyclic groups generated by the elements. Thus we have

$$N \cong \begin{cases} \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/((q-1)/2)\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & \text{if} \quad p = -1 \\ \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/((q-1)/2)\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & \text{if} \quad p = 2 \\ \mathbb{Z}/(p-1)\mathbb{Z} \oplus \mathbb{Z}/((q-1)/2)\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & \text{if} \quad p > 2. \end{cases} \tag{A2.2}$$

In the case B, we define the the subgroup $N$ of $\widetilde{G}$ to be

$$N = \begin{cases} < \widetilde{\sigma}_{-1}, \widetilde{\sigma}_2, \widetilde{\sigma}_q^2 > & \text{if} \quad p = 2 \\ < \widetilde{\sigma}_p, \widetilde{\sigma}_q^2 > & \text{if} \quad p \neq 2 \text{ and } |\widetilde{\sigma}_q| = 2|\sigma_q| \\ < \widetilde{\sigma}_p^2, \widetilde{\sigma}_q > & \text{if} \quad |\widetilde{\sigma}_p| = 2|\sigma_p|. \end{cases} \tag{B2.1}$$

Again $N$ is abelian and has index 2 in $\widetilde{G}$. In addition, we have

$$N \cong \begin{cases} \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} & \text{if} \quad (p,q) = (-1,2) \\ \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/(q-1)\mathbb{Z} & \text{if} \quad p = -1, \ q > 2 \\ \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/(q-1)\mathbb{Z} & \text{if} \quad p = 2 \\ \mathbb{Z}/(p-1)\mathbb{Z} \oplus \mathbb{Z}/(q-1)\mathbb{Z} & \text{if} \quad p > 2. \end{cases} \tag{B2.2}$$

In the case C, we must have $p, q$ are odd prime numbers. Let $v_2(p-1)$ denote the power of 2 in $p-1$. We define the subgroup $N$ of $\widetilde{G}$ to be

$$N = \begin{cases} < \widetilde{\sigma}_p^2\, , \widetilde{\sigma}_q > & \text{if} \quad v_2(p-1) \leq v_2(q-1) \\ < \widetilde{\sigma}_p\, , \widetilde{\sigma}_q^2 > & \text{if} \quad v_2(p-1) > v_2(q-1). \end{cases} \tag{C2.1}$$

Then $N$ is an abelian subgroup of $\widetilde{G}$. When $v_2(p-1) \le v_2(q-1)$, we have

$$|N| = \frac{|\widetilde{\sigma}_p^2| \cdot |\widetilde{\sigma}_q|}{| < \widetilde{\sigma}_p^2 > \cap < \widetilde{\sigma}_q > |} = \frac{(p-1)\cdot 2(q-1)}{2}.$$

So $[\widetilde{G} : N] = 2$ and $N$ is a normal subgroup of $\widetilde{G}$. We have the same result when $v_2(p-1) > v_2(q-1)$. The subgroup $< \widetilde{\sigma}_p^2 , \widetilde{\sigma}_q >$ is always an abelian subgroup of $\widetilde{G}$ of index 2. But we can not get all irreducible representations from the inducement of the representations of this subgroup when $v_2(p-1) > v_2(q-1)$. So we define $N$ in two cases.

Next we determine the structure of the subgroup $N$ in the case C. We consider the case $v_2(p-1) \le v_2(q-1)$ in detail. Let $d = \gcd(\frac{p-1}{2}, q-1)$, $s = (p-1)/2d$ and $t = (q-1)/d$. Choose $u, v \in \mathbb{Z}$ such that $us + vt = 1$. We have the relations

$$(\widetilde{\sigma}_p^2)^{p-1} = 1, \quad (\widetilde{\sigma}_p^2)^{\frac{p-1}{2}} = \varepsilon = \widetilde{\sigma}_q^{q-1}.$$

Let $M$ be the free abelian group generated by two words $\alpha$ , $\beta$. Let

$$\alpha_1 = (p-1)\alpha \ \ ; \ \ \beta_1 = \frac{p-1}{2}\alpha - (q-1)\beta \ ;$$

and let $M_1$ be the subgroup of $M$ generated by $\alpha_1 , \beta_1$. Then $M_1$ is the kernel of the homomorphism

$$M \longrightarrow N; \ \ \alpha \mapsto \widetilde{\sigma}_p^2, \ \beta \mapsto \widetilde{\sigma}_q \ \ .$$

So we have $N \cong M/M_1$. Define the matrix

$$A := \begin{pmatrix} p-1 & \frac{p-1}{2} \\ 0 & 1-q \end{pmatrix} \ .$$

Then $(\alpha_1 , \beta_1) = (\alpha , \beta) \cdot A$. We determine the structure of $M_1$ by considering the standard form of $A$. Define

$$P := \begin{pmatrix} u & v \\ -t & s \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \ ; \qquad Q := \begin{pmatrix} 1 & 2tv-1 \\ -1 & -2tv+2 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}).$$

Then

$$B := PAQ = \begin{pmatrix} d & 0 \\ 0 & -2s(q-1) \end{pmatrix}$$

is the standard form of $A$. Let

$$(\tau , \mu) = (\alpha , \beta)P^{-1} \quad \text{and} \quad (\tau_1 , \mu_1) = (\alpha_1 , \beta_1)Q.$$

Then $(\tau_1\,,\mu_1) = (\tau\,,\mu)B$, $M = \mathbb{Z}\tau \oplus \mathbb{Z}\mu$ and $M_1 = \mathbb{Z}d\tau \oplus \mathbb{Z}2s(q-1)\mu$. We thus have

$$N \cong M/M_1 \cong \mathbb{Z}/d\mathbb{Z} \oplus \mathbb{Z}/2s(q-1)\mathbb{Z}\ .$$

By abuse of notation, we also write

$$(\tau\,,\mu) = (\widetilde{\sigma}_p^2\,, \widetilde{\sigma}_q)P^{-1} = (\widetilde{\sigma}_p^{2s}\widetilde{\sigma}_q^t\,,\ \widetilde{\sigma}_p^{-2v}\widetilde{\sigma}_q^u)\ .$$

Then $\tau\,,\mu$ are of order $d\,, 2s(q-1)$ respectively, and $N$ is a direct sum of $<\tau>$ and $<\mu>$. We have $\widetilde{\sigma}_p^2 = \tau^u\mu^{-t}$ and $\widetilde{\sigma}_q = \tau^v\mu^s$. When $v_2(p-1) > v_2(q-1)$, we get the structure of $N$ in the same way. So in the case (C) we have

$$N \cong \begin{cases} \mathbb{Z}/d\mathbb{Z} \oplus \mathbb{Z}/2s(q-1)\mathbb{Z} & \text{if} \quad v_2(p-1) \le v_2(q-1) \\ \mathbb{Z}/d'\mathbb{Z} \oplus \mathbb{Z}/2s'(p-1)\mathbb{Z} & \text{if} \quad v_2(p-1) > v_2(q-1), \end{cases} \qquad (C2.2)$$

where $d = \gcd(\frac{p-1}{2}, q-1), s = (p-1)/2d$ and $d' = \gcd(p-1, \frac{q-1}{2}), s' = (q-1)/2d'$.

Now we summarize our results in the following

**Proposition 2.1.** *The abelian subgroup $N$ of the group $\widetilde{G}$ of index 2 defined in (A2.1), (B2.1) and (C2.1) has the structure described in (A2.2), (B2.2) and (C2.2) in the cases (A), (B) and (C), respectively. In particular, every irreducible representation of $\widetilde{G}$ has dimension 1 or 2.*

# 3 2-dimensional representations

We determine all irreducible representations of $\widetilde{G}$ in this section. We will use some basic facts from representation theory freely. For the details, see [6].

It is well-known that the 1-dimensional representations of $\widetilde{G}$ correspond bijectively to those of the maximal abelian quotient $G$ of $\widetilde{G}$, which are Dirichlet characters. So we mainly construct the 2-dimensional irreducible representations of $\widetilde{G}$. From the dimension formula of all irreducible representations, we see that $\widetilde{G}$ has $|G|/4$ irreducible representations of dimension 2, up to isomorphism. Let $N$ be the subgroup of $\widetilde{G}$ defined in last section. Let $\widetilde{G} = N \cup \sigma N$ be a decomposition of cosets. If $\rho : N \to \mathbb{C}^*$ is a representation of $N$, the induced representation $\widetilde{\rho}$ of $\rho$ is a representation of $\widetilde{G}$ of dimension 2. The space of the representation $\widetilde{\rho}$ is $V = \mathrm{Ind}_N^{\widetilde{G}}(\mathbb{C}) = \mathbb{C}[\widetilde{G}] \otimes_{\mathbb{C}[N]} \mathbb{C}$ with basis $e_1 = 1 \otimes 1$ and $e_2 = \sigma \otimes 1$. The group homomorphism

$$\widetilde{\rho} : \widetilde{G} \longrightarrow \mathrm{GL}(V) \simeq \mathrm{GL}_2(\mathbb{C})$$

is given under the basis by

$$\widetilde{\rho}(\widetilde{\sigma}) = \begin{pmatrix} \rho(\widetilde{\sigma}) & \rho(\widetilde{\sigma}\sigma) \\ \rho(\sigma^{-1}\widetilde{\sigma}) & \rho(\sigma^{-1}\widetilde{\sigma}\sigma) \end{pmatrix}, \qquad \forall\, \widetilde{\sigma} \in \widetilde{G}, \qquad (3.1)$$

where $\rho(\widetilde{\sigma}) = 0$ if $\widetilde{\sigma} \notin N$. The representation $\widetilde{\rho}$ is irreducible if and only if $\rho \ncong \rho^\tau$ for every $\tau \in \widetilde{G} \setminus N$, where $\rho^\tau$ is the conjugate representation of $\rho$ defined by

$$\rho^\tau(x) = \rho(\tau^{-1} x \tau) \ , \ \ \forall\, x \in N\,.$$

Since $N$ is abelian, we only need to check $\rho \ncong \rho^\sigma$.

Now we begin to construct all 2-dimensional irreducible representations of $\widetilde{G}$. As in last section, we consider the three cases separately. In addition, we consider the case when $p$ and $q$ are odd prime numbers in details, and only state the results in the cases when $p = -1$ or 2.

**3.1. Case A.** Assume $p > 2$. We have in this case $N = \langle \widetilde{\sigma}_p\, , \, \widetilde{\sigma}_q^2\, , \varepsilon \rangle$ and

$$N \cong \mathbb{Z}/(p-1)\mathbb{Z} \oplus \mathbb{Z}/((q-1)/2)\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}\,.$$

Every irreducible representation of $N$ can be written as $\rho_{ijk} : N \longrightarrow \mathbb{C}^*$ with

$$\rho_{ijk}(\widetilde{\sigma}_p) = \zeta_{p-1}^i\ ;\ \rho_{ijk}(\widetilde{\sigma}_q^2) = \zeta_{q-1}^{2j}\ ;\ \rho_{ijk}(\varepsilon) = (-1)^k\,.$$

where $0 \le i < p-1$, $0 \le j < \frac{q-1}{2}$ and $k = 0, 1$. Since $\widetilde{G} = N \cup \widetilde{\sigma}_q N$ and $\rho_{ijk}^{\widetilde{\sigma}_q}(\widetilde{\sigma}_p) = \rho_{ijk}(\varepsilon)\rho_{ijk}(\widetilde{\sigma}_p) = (-1)^k \rho_{ijk}(\widetilde{\sigma}_p)$, we have

$$\rho_{ijk}^{\widetilde{\sigma}_q} \ncong \rho_{ijk} \Longleftrightarrow k = 1.$$

Write $\rho_{ij} = \rho_{ij1}$. The induced representation $\widetilde{\rho}_{ij} : \widetilde{G} \longrightarrow \mathrm{GL}_2(\mathbb{C})$ of $\rho_{ij}$ is given by

$$\widetilde{\rho}_{ij}(\widetilde{\sigma}_p) = \begin{pmatrix} \zeta_{p-1}^i & 0 \\ 0 & -\zeta_{p-1}^i \end{pmatrix},\ \widetilde{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^{2j} \\ 1 & 0 \end{pmatrix},\ \widetilde{\rho}_{ij}(\varepsilon) = -I, \qquad (A3.1)$$

where $I$ is the identity matrix of degree 2. Since

$$\widetilde{\rho}_{ij}(\widetilde{\sigma}_p^2) = \begin{pmatrix} \zeta_{p-1}^{2i} & 0 \\ 0 & \zeta_{p-1}^{2i} \end{pmatrix} \quad \text{and} \quad \widetilde{\rho}_{ij}(\widetilde{\sigma}_q^2) = \begin{pmatrix} \zeta_{q-1}^{2j} & 0 \\ 0 & \zeta_{q-1}^{2j} \end{pmatrix},$$

we see that the representations $\widetilde{\rho}_{ij}$ with $0 \le i < \frac{p-1}{2}$, $0 \le j < \frac{q-1}{2}$ are irreducible and are not isomorphic to each other, by considering the values of the characters of these representations at $\widetilde{\sigma}_p^2$ and $\widetilde{\sigma}_q^2$. The number of these representations is $\frac{p-1}{2} \cdot \frac{q-1}{2} = \frac{|G|}{4}$. So they are all the irreducible representations of $\widetilde{G}$ of dimension 2.

Similarly, when $p = -1$, all irreducible representations of $\widetilde{G}$ of dimension 2 are $\widetilde{\rho}_j$ with $0 \le j < \frac{q-1}{2}$, where

$$\widetilde{\rho}_j(\widetilde{\sigma}_{-1}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \widetilde{\rho}_j(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^{2j} \\ 1 & 0 \end{pmatrix}, \quad \widetilde{\rho}(\varepsilon) = -I \qquad (A3.2)$$

and when $p = 2$, all irreducible representations of $\widetilde{G}$ of dimension 2 are $\bar{\rho}_{ij}$ with $0 \leq i \leq 1$ and $0 \leq j < \frac{q-1}{2}$, where $\bar{\rho}_{ij}(\varepsilon) = -I$ and

$$\bar{\rho}_{ij}(\widetilde{\sigma}_{-1}) = (-1)^i I, \ \bar{\rho}_{ij}(\widetilde{\sigma}_2) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \ \bar{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^{2j} \\ 1 & 0 \end{pmatrix}. \qquad (A3.3)$$

**3.2 Case B.** Assume $p > 2$ and $|\widetilde{\sigma}_q| = 2|\sigma_q|$. Then $N = \langle \widetilde{\sigma}_p \, , \, \widetilde{\sigma}_q^2 \rangle$, and

$$N \cong \mathbb{Z}/(p-1)\mathbb{Z} \oplus \mathbb{Z}/(q-1)\mathbb{Z}.$$

Any irreducible representation of $N$ has the form $\rho_{ij} : N \longrightarrow \mathbb{C}^*$, where

$$\rho_{ij}(\widetilde{\sigma}_p) = \zeta_{p-1}^i \ , \ \rho_{ij}(\widetilde{\sigma}_q^2) = \zeta_{q-1}^j \ , \ \rho_{ij}(\varepsilon) = \rho_{ij}(\widetilde{\sigma}_q^2)^{\frac{q-1}{2}} = (-1)^j \ ,$$

and $0 \leq i < p-1, \ 0 \leq j < q-1$. It is easy to check that

$$\rho_{ij}^{\widetilde{\sigma}_q} \not\cong \rho_{ij} \Longleftrightarrow j \equiv 1 \pmod 2 \ .$$

The induced representation $\widetilde{\rho}_{ij} : \widetilde{G} \longrightarrow \mathrm{GL}_2(\mathbb{C})$ of $\rho_{ij}$ with odd $j$ is given by

$$\widetilde{\rho}_{ij}(\widetilde{\sigma}_p) = \begin{pmatrix} \zeta_{p-1}^i & 0 \\ 0 & -\zeta_{p-1}^i \end{pmatrix}, \ \ \widetilde{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^j \\ 1 & 0 \end{pmatrix}. \qquad (B3.1)$$

Since

$$\widetilde{\rho}_{ij}(\widetilde{\sigma}_p^2) = \begin{pmatrix} \zeta_{p-1}^{2i} & 0 \\ 0 & \zeta_{p-1}^{2i} \end{pmatrix} \quad \text{and} \quad \widetilde{\rho}_{ij}(\widetilde{\sigma}_q^2) = \begin{pmatrix} \zeta_{q-1}^j & 0 \\ 0 & \zeta_{q-1}^j \end{pmatrix},$$

we see that the representations $\widetilde{\rho}_{ij}$ with $0 \leq i < \frac{p-1}{2}$ and $0 \leq j < q-1, \ 2 \nmid j$ are irreducible and are not isomorphic to each other. The number of these representations is $\frac{|G|}{4}$. So they are all the irreducible representations of $\widetilde{G}$ of dimension 2.

Similarly, when $(p, q) = (-1, 2)$, there is only one irreducible representation $\widetilde{\rho}_0$ of dimension 2 defined by

$$\widetilde{\rho}_0(\widetilde{\sigma}_{-1}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \text{and} \quad \widetilde{\rho}_0(\widetilde{\sigma}_2) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \qquad (B3.2)$$

When $p = -1$ and $q > 2$, all irreducible representations of dimension 2 are $\widetilde{\rho}_j$ with $0 \leq j < q-1, \ 2 \nmid j$, where $\widetilde{\rho}_j$ is defined by

$$\widetilde{\rho}_j(\widetilde{\sigma}_{-1}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad \widetilde{\rho}_j(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^j \\ 1 & 0 \end{pmatrix}. \qquad (B3.3)$$

When $p = 2$, all irreducible representations of dimension 2 are $\bar{\rho}_{ij}$ with $0 \leq i \leq 1$ and $0 \leq j < q-1, \ 2 \nmid j$, where $\bar{\rho}_{ij}$ is defined by

$$\bar{\rho}_{ij}(\widetilde{\sigma}_{-1}) = (-1)^i I, \ \bar{\rho}_{ij}(\widetilde{\sigma}_2) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \ \bar{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^j \\ 1 & 0 \end{pmatrix}. \qquad (B3.4)$$

When $|\widetilde{\sigma}_p| = 2|\sigma_p|$, all irreducible representations of dimension 2 are $\hat{\rho}_{ij}$ with $0 \le i < p-1,\ 2 \nmid i$ and $0 \le j < \frac{q-1}{2}$, where $\hat{\rho}_{ij}$ is defined by

$$\hat{\rho}_{ij}(\widetilde{\sigma}_p) = \begin{pmatrix} 0 & \zeta_{p-1}^i \\ 1 & 0 \end{pmatrix}, \quad \hat{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} \zeta_{q-1}^j & 0 \\ 0 & -\zeta_{q-1}^j \end{pmatrix}. \qquad (B3.5)$$

**3.3. Case C.** Assume $v_2(p-1) \le v_2(q-1)$. Let

$$d = \gcd(\frac{p-1}{2}, q-1)\ ,\ s = \frac{p-1}{2d}, t = \frac{q-1}{d}\ ;\ \ us + vt = 1$$

as before. We must have that $t$ is even and $u$ is odd. Let $\tau = \widetilde{\sigma}_p^{2s} \cdot \widetilde{\sigma}_q^t$ and $\mu = \widetilde{\sigma}_p^{-2v} \cdot \widetilde{\sigma}_q^u$. Then $N = \langle \widetilde{\sigma}_p^2\, , \widetilde{\sigma}_q \rangle = \langle \tau\, , \mu \rangle$ and

$$N \cong \mathbb{Z}/d\mathbb{Z} \oplus \mathbb{Z}/2s(q-1)\mathbb{Z}.$$

Any irreducible representation $\rho_{ij} : N \longrightarrow \mathbb{C}^*$ is of the form

$$\rho_{ij}(\tau) = \zeta_d^i = \zeta_{(p-1)(q-1)}^{2s(q-1)i} \quad \text{and} \quad \rho_{ij}(\mu) = \zeta_{2s(q-1)}^j = \zeta_{(p-1)(q-1)}^{dj}.$$

From $\widetilde{\sigma}_p^2 = \tau^u \mu^{-t}$ and $\widetilde{\sigma}_q = \tau^v \mu^s$, we have

$$\rho_{ij}(\widetilde{\sigma}_p^2) = \zeta_{p-1}^{2sui-j}\ ;\ \rho_{ij}(\widetilde{\sigma}_q) = \zeta_{2(q-1)}^{2tvi+j}\ ;\ \rho_{ij}(\varepsilon) = \rho_{ij}(\widetilde{\sigma}_p^2)^{\frac{p-1}{2}} = (-1)^j\ .$$

It is easy to show

$$\rho_{ij}^{\widetilde{\sigma}_p} \not\cong \rho_{ij} \Longleftrightarrow j \equiv 1 \pmod 2\ .$$

The induced representation $\widetilde{\rho}_{ij} : \widetilde{G} \longrightarrow \mathrm{GL}_2(\mathbb{C})$ of $\rho_{ij}$ with odd $j$ is given by

$$\widetilde{\rho}_{ij}(\tau) = \begin{pmatrix} \zeta_d^i & 0 \\ 0 & \zeta_d^i \end{pmatrix}\ ; \quad \widetilde{\rho}_{ij}(\mu) = \begin{pmatrix} \zeta_{2s(q-1)}^j & 0 \\ 0 & -\zeta_{2s(q-1)}^j \end{pmatrix}.$$

Here in the first equality we used the fact that $t$ is even, and in the second equality we used the fact that $u$ is odd. Furthermore we have

$$\widetilde{\rho}_{ij}(\widetilde{\sigma}_p) = \begin{pmatrix} 0 & \zeta_{p-1}^{2sui-j} \\ 1 & 0 \end{pmatrix}\ ;\ \widetilde{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} \zeta_{2(q-1)}^{2tvi+j} & 0 \\ 0 & -\zeta_{2(q-1)}^{2tvi+j} \end{pmatrix}. \qquad (C3.1)$$

By considering the values of the character of $\widetilde{\rho}_{ij}$ at $\tau$ and $\mu^2$, we see that all the representations $\widetilde{\rho}_{ij}$ with $0 \le i < d$ and $0 \le j < s(q-1),\ 2 \nmid j$ are irreducible and are not isomorphic to each other. The number of these representations is $d \cdot \frac{s(q-1)}{2} = \frac{|G|}{4}$. So they are all the irreducible representations of $\widetilde{G}$ of dimension 2.

Similarly, if $v_2(p-1) > v_2(q-1)$, we let

$$d' = \gcd(p-1\,, \frac{q-1}{2})\ ,\ s' = \frac{p-1}{d}\,, t' = \frac{q-1}{2d}\ ;\ u's' + v't' = 1.$$

Then all the irreducible representations of $\widetilde{G}$ of dimension 2 are $\hat{\rho}_{ij}$ with $0 \le i < d'$ and $0 \le j < t'(p-1),\ 2 \nmid j$, where $\hat{\rho}_{ij}$ is defined by

$$\hat{\rho}_{ij}(\widetilde{\sigma}_p) = \begin{pmatrix} \zeta_{2(p-1)}^{2s'u'i+j} & 0 \\ 0 & -\zeta_{2(p-1)}^{2s'u'i+j} \end{pmatrix}\ ;\ \hat{\rho}_{ij}(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^{2t'v'i-j} \\ 1 & 0 \end{pmatrix}\ . \qquad (C3.2)$$

Let $\mathrm{R}^2(\widetilde{G})$ be the set of all irreducible representations, up to isomorphism, of $\widetilde{G}$ of dimension 2. As a summary, we have proved the following.

**Theorem 3.1.** *All 2-dimensional irreducible representations of $\widetilde{G}$ are induced from the representations of $N$. In detail, we have*

*In the case (A)*

$$\mathrm{R}^2(\widetilde{G}) = \begin{cases} \{\widetilde{\rho}_j \mid 0 \le j < \frac{q-1}{2}\} & \text{if } p = -1 \\ \{\bar{\rho}_{ij} \mid i = 0, 1,\ 0 \le j < \frac{q-1}{2}\} & \text{if } p = 2 \\ \{\widetilde{\rho}_{ij} \mid 0 \le i < \frac{p-1}{2},\ 0 \le j < \frac{q-1}{2}\} & \text{if } p > 2, \end{cases}$$

*where $\widetilde{\rho}_j$, $\bar{\rho}_{ij}$ and $\widetilde{\rho}_{ij}$ are defined in (A3.2), (A3.3) and (A3.1) respectively.*

*In the case (B)*

$$\mathrm{R}^2(\widetilde{G}) = \begin{cases} \{\widetilde{\rho}_0\} & \text{if } (p,q) = (-1,2) \\ \{\widetilde{\rho}_j \mid 0 \le j < q-1,\ 2 \nmid j\} & \text{if } p = -1,\ q > 2 \\ \{\bar{\rho}_{ij} \mid i = 0, 1,\ 0 \le j < q-1,\ 2 \nmid j\} & \text{if } p = 2 \\ \{\hat{\rho}_{ij} \mid 0 \le i < p-1,\ 2 \nmid i,\ 0 \le j < \frac{q-1}{2}\} & \text{if } |\widetilde{\sigma}_p| = 2|\sigma_p| \\ \{\widetilde{\rho}_{ij} \mid 0 \le i < \frac{p-1}{2},\ 0 \le j < q-1,\ 2 \nmid j\} & \text{otherwise,} \end{cases}$$

*where $\widetilde{\rho}_0$, $\widetilde{\rho}_j$, $\bar{\rho}_{ij}$, $\hat{\rho}_{ij}$ and $\widetilde{\rho}_{ij}$ are defined in (B3.2), (B3.3), (B3.4), (B3.5) and (B3.1) respectively.*

*In the case (C)*

$$\mathrm{R}^2(\widetilde{G}) = \begin{cases} \{\widetilde{\rho}_{ij} \mid 0 \le i < d,\ 0 \le j < s(q-1),\ 2 \nmid j\} & \text{if } v_2(p-1) \le v_2(q-1), \\ \{\hat{\rho}_{ij} \mid 0 \le i < d',\ 0 \le j < t'(p-1),\ 2 \nmid j\} & \text{otherwise,} \end{cases}$$

*where $\widetilde{\rho}_{ij}$ and $\hat{\rho}_{ij}$ are defined in (C3.1) and (C3.2) respectively.*

# 4 The Frobenious maps

This section is a preparation for the next section to compute the Artin $L$-functions of the quasi-cyclotomic fields $\widetilde{K}$ when $p = -1$. For a prime number $\ell$ which is unramified in $\widetilde{K}/K$, let $I_\ell$ (resp. $\widetilde{I_\ell}$) be the inertia group of $\ell$ in the extension $K/\mathbb{Q}$ (resp. $\widetilde{K}/\mathbb{Q}$). Let $\mathrm{Fr}_\ell$ be the Frobenious automorphism of $\ell$ in $G/I_\ell$ and $\widetilde{\mathrm{Fr}}_\ell$ the Frobenious automorphism of $\ell$ in $\widetilde{G}/\widetilde{I_\ell}$ associated to some prime ideal over $\ell$. In this section we determine $\widetilde{\mathrm{Fr}}_\ell$ by $\mathrm{Fr}_\ell$ for $\ell = 2$.

From now on, we always assume that $p = -1$, namely, $K = \mathbb{Q}(\zeta_{4q})$ and $\widetilde{K} = K(\sqrt[4]{q^*})$. For a prime number $\ell$, we say that $\ell$ is ramified (resp. inertia, splitting) in the relative quadratic extension $\widetilde{K}/K$ if the prime ideals of $K$ over $\ell$ are ramified (resp. inertia, splitting) in $\widetilde{K}$. In [Sect.5, 7] we have determined the decomposition nature of odd prime numbers in $\widetilde{K}/K$. Now we determine the decomposition nature of 2 in $\widetilde{K}/K$.

**Proposition 4.1.** *If $q = 2$, then 2 is ramified in $\widetilde{K}/K$. If $q$ is odd, then 2 is unramified in $\widetilde{K}/K$ if and only if $(\frac{2}{q}) = 1$, and in this case 2 splits in $\widetilde{K}/K$ if $q^* \equiv 1\mathrm{mod}16$ and is inertia in $\widetilde{K}/K$ if $q^* \equiv 1\mathrm{mod}8$ but $q^* \not\equiv 1\mathrm{mod}16$.*

*Proof.* We first consider the case $q = 2$. The unique prime ideal of $K$ over 2 is the principal ideal generated by $\pi := 1 - \zeta_8$. Since the ramification degree of 2 in $K/\mathbb{Q}$ is 4 and $\sqrt{2} = \pi(\pi + 2\zeta_8)\zeta_8$, we have that 2 is ramified in $\widetilde{K}/K$ if and only if $x^2 \equiv \sqrt{2}\mathrm{mod}\pi^{10}$ is not solvable in the ring $O_K$ of the integers of $K$ by [2], which is equivalent to that $(1 + \frac{2}{\pi}\zeta_8)\zeta_8\mathrm{mod}\pi^8$ is not a square. Since $2 = u\pi^4$ for some unit $u$, we have

$$(1 + \frac{2}{\pi}\zeta_8)\zeta_8 \equiv \zeta_8 \equiv (1 - \pi)\mathrm{mod}\pi^3,$$

namely $(1 + \frac{2}{\pi}\zeta_8)\zeta_8\mathrm{mod}\pi^3$ is not a square. So 2 is ramified in $\widetilde{K}/K$.

Now we assume that $q$ is odd. Let $\pi_2 = 1 - \zeta_4$. Then 2 is unramified in $\widetilde{K}/K$ if and only if $x^2 \equiv \sqrt{q^*}\mathrm{mod}\pi_2^4$ is solvable in $O_K$. Furthermore, 2 splits in $\widetilde{K}/K$ if and only if $x^2 \equiv \sqrt{q^*}\mathrm{mod}\pi_2^5$ is solvable in $O_K$. By Gauss sum we have

$$\sqrt{q^*} = \sum_{a=1}^{q-1} (\frac{a}{q})\zeta_q^a = 1 + 2 \sum_{(\frac{a}{q})=1} \zeta_q^a.$$

Let $\alpha = \sum_{(\frac{a}{q})=1} \zeta_q^a$, $\beta = \sum_{(\frac{a}{q})=1} \zeta_{2q}^a$, and $\gamma = \sum_{(\frac{a}{q})=1} \sum_{(\frac{b}{q})=1,\ a<b} \zeta_{2q}^{a+b}$, where in the summations $a, b$ run over $1, 2, \cdots, q-1$. Then $\alpha = \beta^2 - 2\gamma$, from which and the equality $2 = \pi_2^2 - \pi_2^3$, we have

$$\begin{aligned}\sqrt{q^*} &= 1 + 2\beta^2 - 4\gamma = 1 + \pi_2^2\beta^2 - \pi_2^3\beta^2 - 4\gamma \\ &\equiv (1 + \pi_2\beta)^2 - \pi_2^3(\beta + \beta^2) + \pi_2^4(\beta - \gamma) \\ &\equiv (1 + \pi_2\beta)^2 - \pi_2^3(\alpha + \beta) + \pi_2^4(\beta + \gamma)\mathrm{mod}\pi_2^5.\end{aligned}$$

Since $\zeta_{2q} = -\zeta_q^{-\frac{q-1}{2}} = -\zeta_q^t$, where $t$ is the inverse of 2 in $(\mathbb{Z}/q\mathbb{Z})^*$, we see $\beta = \sum_{(\frac{a}{q})=1}(-1)^a\zeta_q^{ta} \equiv \sum_{(\frac{a}{q})=1}\zeta_q^{ta}\mathrm{mod}2$. So if $(\frac{2}{q}) = 1$ we have $\alpha \equiv \beta\mathrm{mod}2$ and thus 2 is unramified in $\widetilde{K}/K$, and if $(\frac{2}{q}) = -1$ we have $\alpha+\beta \equiv \sum_{a=1}^{q-1}\zeta_q^a = -1\mathrm{mod}2$ and thus 2 is ramified in $\widetilde{K}/K$.

Now we assume $(\frac{2}{q}) = 1$. Then $\sqrt{q^*}\mathrm{mod}\pi_2^5$ is a square if and only if $\pi_2 \mid \beta+\gamma$. We consider $2(\beta+\gamma)$. Since $\alpha \equiv \beta\mathrm{mod}2$, we have

$$2(\beta+\gamma) = 2\beta+\beta^2-\alpha \equiv \alpha(\alpha+1)\mathrm{mod}4.$$

From $\sqrt{q^*} = 1+2\alpha$, we see $\alpha(\alpha+1) = \frac{q^*-1}{4}$. Since $8 \mid q^*-1$ under the assumption $(\frac{2}{q}) = 1$, we have $\beta+\gamma \equiv \frac{q^*-1}{8}\mathrm{mod}2$. So $\pi_2 \mid \beta+\gamma$ if and only if $\pi_2 \mid \frac{q^*-1}{8}$, namely $2 \mid \frac{q^*-1}{8}$. We complete the proof. □

By the way, we have determined the ring $O_{\widetilde{K}}$ of the integers of $\widetilde{K}$. In fact, we have

**Corollary 4.2.** *Assume that $q$ is an odd prime number. Let $t_q = (q-1)/4$ if $q \equiv 1\mathrm{mod}4$, and $t_q = (q-3)/4$ if $q \equiv 3\mathrm{mod}4$. Then*

$$O_{\widetilde{K}} = \begin{cases} \mathbb{Z}\left[\zeta_{4q}, \frac{\sqrt[4]{q^*}+1+\pi_2\beta}{\pi_2(\sin[\frac{1}{q}])^{t_q}}\right] & \text{if } (\frac{2}{q}) = -1 \\ \mathbb{Z}\left[\zeta_{4q}, \frac{\sqrt[4]{q^*}+1+\pi_2\beta}{2(\sin[\frac{1}{q}])^{t_q}}\right] & \text{if } (\frac{2}{q}) = 1. \end{cases}$$

*Proof.* See [Th.2, 7]. □

Now we assume that 2 is unramified in $\widetilde{K}/K$. Let $\mathrm{Fr}_2 \in G$ such that $\mathrm{Fr}_2(\zeta_4) = 1$ and $\mathrm{Fr}_2(\zeta_q) = \zeta_q^2$. It is a Frobenious element of 2 in $G$ modulo $I_2$. We have $\mathrm{Fr}_2 = \sigma_2^{b_2}$ with $2 \mid b_2$ for $(\frac{2}{q}) = 1$. Thus $\widetilde{\mathrm{Fr}}_2 = \widetilde{\sigma}_2^{b_2}$ or $\widetilde{\mathrm{Fr}}_2 = \widetilde{\sigma}_2^{b_2}\varepsilon$. We need to determine $\widetilde{\mathrm{Fr}}_2$ completely. Since $(\frac{2}{q}) = 1$, we have

$$\sqrt{q^*} \equiv (1+\pi_2\alpha)^2 + \pi_2^4(\beta+\gamma)\mathrm{mod}\pi_2^5.$$

Write $u = 1+\pi_2\alpha$ for simplicity. Since $\sqrt{q^*} \equiv u^2\mathrm{mod}\pi_2^4$, we see $\frac{\sqrt[4]{q^*}-u}{2} \in O_{\widetilde{K}}$. Let $\wp$ be the prime ideal of $\widetilde{K}$ over 2 associated to $\widetilde{\mathrm{Fr}}_2$. By the definition, we have

$$\widetilde{\mathrm{Fr}}_2\left(\frac{\sqrt[4]{q^*}-u}{2}\right) \equiv \left(\frac{\sqrt[4]{q^*}-u}{2}\right)^2 \equiv (\beta+\gamma) + \frac{\sqrt[4]{q^*}-u}{2}\mathrm{mod}\wp.$$

On the other hand, since $\widetilde{\sigma}_q^{b_2}(\sqrt[4]{q^*}) = (-1)^{\frac{b_2}{2}}\sqrt[4]{q^*}$ and $\widetilde{\sigma}_q^{b_2}(u) = u$ for $2 \mid b_2$, we have

$$\widetilde{\sigma}_q^{b_2}(\frac{\sqrt[4]{q^*}-u}{2}) = \frac{(-1)^{\frac{b_2}{2}}\sqrt[4]{q^*}-u}{2}$$

and

$$\widetilde{\sigma}_q^{b_2}\varepsilon(\frac{\sqrt[4]{q^*}-u}{2})=\frac{(-1)^{\frac{b_2}{2}+1}\sqrt[4]{q^*}-u}{2}.$$

So if $2\mid\frac{b_2}{2}$ we have $\widetilde{\mathrm{Fr}}_2=\widetilde{\sigma}_2^{b_2}$ if and only if $\pi_2\mid\beta+\gamma$ (namely 2 splits in $\widetilde{K}/K$), and if $2\nmid\frac{b_2}{2}$ we have $\widetilde{\mathrm{Fr}}_2=\widetilde{\sigma}_2^{b_2}$ if and only if $\pi_2\nmid\beta+\gamma$ (namely 2 is inertia in $\widetilde{K}/K$). In the case $q\equiv 3\mathrm{mod}4$, we can always assume that $2\nmid\frac{b_2}{2}$, since if $4\mid b_2$, we may replace $b_2$ by $b_2+(q-1)$. In the case $q\equiv 1\mathrm{mod}4$, we have $2\mid\frac{b_2}{2}\iff 2^{\frac{q-1}{4}}\equiv 1\mathrm{mod}q\iff q$ has the form $A^2+64B^2$ for $A,B\in\mathbb{Z}$, by the Exercise 28 in Chap.5 in [5]. So we get the following result

**Proposition 4.3.** *Assume that 2 is unramified in $\widetilde{K}/K$. Let $\mathrm{Fr}_2=\sigma_2^{b_2}$. We have $2\mid b_2$. If $q\equiv 3\mathrm{mod}4$, we always assume $b_2\equiv 2\mathrm{mod}4$. Let $P_0$ be the set of the prime numbers of the form $A^2+64B^2$ with $A,B\in\mathbb{Z}$. Then we have*

$$\widetilde{\mathrm{Fr}}_2=\begin{cases}\widetilde{\sigma}_2^{b_2} & \text{if}\quad q\notin P_0, 16\nmid q^*-1,\ \text{or}\ q\in P_0, 16\mid q^*-1\\ \widetilde{\sigma}_2^{b_2}\varepsilon & \text{if}\quad q\in P_0, 16\nmid q^*-1,\ \text{or}\ q\notin P_0, 16\mid q^*-1.\end{cases}$$

The following lemma is useful in next section.

**Lemma 4.4.** *We have $\varepsilon\in\widetilde{I}_\ell$ if and only if $\ell$ is ramified in $\widetilde{K}/K$.*

*Proof.* The canonical projection $\widetilde{G}\longrightarrow G\simeq\widetilde{G}/\langle\varepsilon\rangle$ induces a surjective homomorphism $\widetilde{I}_\ell\longrightarrow I_\ell$ which implies the isomorphism $\widetilde{I}_\ell/<\varepsilon>\cap\widetilde{I}_\ell\cong I_\ell$. Thus $\ell$ is ramified in $\widetilde{K}/K\iff|\widetilde{I}_\ell|=2|I_\ell|\iff|\widetilde{I}_\ell\cap<\varepsilon>|=2\iff\varepsilon\in\widetilde{I}_\ell$. $\square$

# 5 The Artin $L$-functions

In this section we compute the Artin $L$-functions of the quasi-cyclotomic fields $\widetilde{K}=\mathbb{Q}(\zeta_{4q},\sqrt[4]{q^*})$.

The $L$-functions associated to the 1-dimensional representations of $\widetilde{G}$ are the well-known Dirichlet $L$-functions. So we mainly compute the $L$-functions associated to the 2-dimensional irreducible representation of $\widetilde{G}$. Let $\varphi:\widetilde{G}\to\mathrm{GL}(V)$ be a 2-dimensional irreducible representations. The Artin $L$-function $L(\varphi,s)$ associated to $\varphi$ is defined as the product of the local factors

$$L(\varphi,s)=\prod_{\ell:\text{prime}}L_\ell(\varphi,s),$$

where the local factors are defined as $L_\ell(\varphi,s)=\det(1-\varphi(\widetilde{\mathrm{Fr}}_\ell)\ell^{-s}|V^{\widetilde{I}_\ell})^{-1}$. Now we begin to compute them. First we notice that if $\ell$ is ramified in $\widetilde{K}/K$, then $V^{\widetilde{I}_\ell}=0$ and $L_\ell(\varphi,s)=1$, which is due to the facts that $\varepsilon\in\widetilde{I}_\ell$ by Lem.4.4 and $\varphi(\varepsilon)=-I$ for any irreducible representation $\varphi$ of $\widetilde{G}$ by Th.3.1.

**5.1 The case $q = 2$.** By section 3, there is only one 2-dimensional representation $\widetilde{\rho}_0$ in this case, which is defined as

$$\widetilde{\rho}_0(\widetilde{\sigma}_{-1}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \text{and} \quad \widetilde{\rho}_0(\widetilde{\sigma}_2) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Since 2 is ramified in $\widetilde{K}/K$, we have $L_2(\widetilde{\rho}_0, s) = 1$. Assume that $\ell$ is an odd prime number.

If $\ell \equiv 7\mathrm{mod}8$, then $\mathrm{Fr}_\ell = \sigma_{-1}$ and thus $\widetilde{\mathrm{Fr}}_\ell = \widetilde{\sigma}_{-1}$ or $\widetilde{\sigma}_{-1}\varepsilon$. In any case we have

$$L_\ell(\widetilde{\rho}_0, s) = \det\left( I \pm \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \ell^{-s} \right)^{-1} = (1 - \ell^{-2s})^{-1}.$$

If $\ell \equiv 5\mathrm{mod}8$, then $\mathrm{Fr}_\ell = \sigma_2$ and thus $\widetilde{\mathrm{Fr}}_\ell = \widetilde{\sigma}_2$ or $\widetilde{\sigma}_2\varepsilon$. We have

$$L_\ell(\widetilde{\rho}_0, s) = \det\left( I \pm \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \ell^{-s} \right)^{-1} = (1 + \ell^{-2s})^{-1}.$$

If $\ell \equiv 3\mathrm{mod}8$, then $\mathrm{Fr}_\ell = \sigma_{-1}\sigma_2$ and thus $\widetilde{\mathrm{Fr}}_\ell = \widetilde{\sigma}_{-1}\widetilde{\sigma}_2$ or $\widetilde{\sigma}_{-1}\widetilde{\sigma}_2\varepsilon$. We have

$$L_\ell(\widetilde{\rho}_0, s) = \det\left( I \pm \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \ell^{-s} \right)^{-1} = (1 - \ell^{-2s})^{-1}.$$

If $\ell \equiv 1\mathrm{mod}8$, then $\mathrm{Fr}_\ell = 1$ and thus $\widetilde{\mathrm{Fr}}_\ell = 1$ or $\varepsilon$. In this case, we must determine $\widetilde{\mathrm{Fr}}_\ell$ completely. Since $\widetilde{\mathrm{Fr}}_\ell(\sqrt[4]{2}) \equiv (\sqrt[4]{2})^\ell \mathrm{mod}\wp$ for the prime ideal $\wp$ of $\widetilde{K}$ over $\ell$ associated to $\widetilde{\mathrm{Fr}}_\ell$, we have $\widetilde{\mathrm{Fr}}_\ell = 1$ if $2^{\frac{\ell-1}{4}} \equiv 1\mathrm{mod}\ell$, and $\widetilde{\mathrm{Fr}}_\ell = \varepsilon$ if $2^{\frac{\ell-1}{4}} \equiv -1\mathrm{mod}\ell$. As in last section, we have that for $\ell \equiv 1\mathrm{mod}8$, $2^{\frac{\ell-1}{4}} \equiv 1\mathrm{mod}\ell$ if and only if $\ell \in P_0$. So we have

$$L_\ell(\widetilde{\rho}_0, s) = \begin{cases} (1 - \ell^{-s})^{-2} & \text{if } \ell \in P_0 \\ (1 + \ell^{-s})^{-2} & \text{otherwise.} \end{cases}$$

We get the Artin $L$-function in the case $(p, q) = (-1, 2)$ as follows.

$$\begin{aligned} L(\widetilde{\rho}_0, s) = & \prod_{\ell \equiv 3 \text{ or } 7(8)} (1 - \ell^{-2s})^{-1} \cdot \prod_{\ell \equiv 5(8)} (1 + \ell^{-2s})^{-1} \\ & \times \prod_{\ell \in P_0} (1 - \ell^{-s})^{-2} \cdot \prod_{\ell \equiv 1(8),\ \ell \notin P_0} (1 + \ell^{-s})^{-2}. \end{aligned} \tag{5.1}$$

**5.2 The case $q$ is odd.** All 2-dimensional irreducible representations of $\widetilde{G}$ are $\widetilde{\rho}_j$ with $0 \le j < q-1, 2 \mid j$ if $q \equiv 1\mathrm{mod}4$, and $0 \le j < q-1, 2 \nmid j$ if $q \equiv 3\mathrm{mod}4$, where $\widetilde{\rho}_j$ is defined by

$$\widetilde{\rho}_j(\widetilde{\sigma}_{-1}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \widetilde{\rho}_j(\widetilde{\sigma}_q) = \begin{pmatrix} 0 & \zeta_{q-1}^j \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \widetilde{\rho}_j(\varepsilon) = -I.$$

We first determine the local factors $L_\ell(\widetilde{\rho}_j, s)$ for $\ell \neq 2, q$. For such $\ell$, we have $V^{\widetilde{I}_\ell} = V$. Let $\mathrm{Fr}_\ell = \sigma_{-1}^{a_\ell}\sigma_q^{b_\ell}$, which is equivalent to $\ell \equiv (-1)^{a_\ell} \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$, where $g$ is the primitive root $\bmod q$ associated to $\sigma_q$. It is easy to compute that

$$\widetilde{\rho}_j(\widetilde{\sigma}_q^{b_\ell}) = \begin{pmatrix} 0 & \zeta_{q-1}^j \\ 1 & 0 \end{pmatrix}^{b_\ell} = \begin{cases} \zeta_{2(q-1)}^{jb_\ell} I & \text{if } 2 \mid b_\ell \\ \begin{pmatrix} 0 & \zeta_{2(q-1)}^{j(b_\ell+1)} \\ \zeta_{2(q-1)}^{j(b_\ell-1)} & 0 \end{pmatrix} & \text{if } 2 \nmid b_\ell. \end{cases}$$

Furthermore, we have

$$\det(I - \widetilde{\rho}_j(\widetilde{\sigma}_{-1}^{a_\ell}\widetilde{\sigma}_q^{b_\ell})\ell^{-s}) = \begin{cases} (1 - \zeta_{2(q-1)}^{jb_\ell}\ell^{-s})^2 & \text{if } a_\ell = 0,\ 2 \mid b_\ell \\ 1 - \zeta_{q-1}^{jb_\ell}\ell^{-2s} & \text{if } a_\ell = 0,\ 2 \nmid b_\ell \\ & \text{or } a_\ell = 1,\ 2 \mid b_\ell \\ 1 + \zeta_{q-1}^{jb_\ell}\ell^{-2s} & \text{if } a_\ell = 1,\ 2 \nmid b_\ell \end{cases}$$

and

$$\det(I + \widetilde{\rho}_j(\widetilde{\sigma}_{-1}^{a_\ell}\widetilde{\sigma}_q^{b_\ell})\ell^{-s}) = \begin{cases} (1 + \zeta_{2(q-1)}^{jb_\ell}\ell^{-s})^2 & \text{if } a_\ell = 0,\ 2 \mid b_\ell \\ 1 - \zeta_{q-1}^{jb_\ell}\ell^{-2s} & \text{if } a_\ell = 0,\ 2 \nmid b_\ell \\ & \text{or } a_\ell = 1,\ 2 \mid b_\ell \\ 1 + \zeta_{q-1}^{jb_\ell}\ell^{-2s} & \text{if } a_\ell = 1,\ 2 \nmid b_\ell. \end{cases}$$

So we get

$$L_\ell(\widetilde{\rho}_j, s) = (1 - \zeta_{q-1}^{jb_\ell}\ell^{-2s})^{-1}$$

if $\ell \equiv 1 \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$ with $2 \nmid b_\ell$, or if $\ell \equiv 3 \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$ with $2 \mid b_\ell$, and

$$L_\ell(\widetilde{\rho}_j, s) = (1 + \zeta_{q-1}^{jb_\ell}\ell^{-2s})^{-1}$$

if $\ell \equiv 3 \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$ with $2 \nmid b_\ell$.

To compute the local factors when $\ell \equiv 1 \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$ with $2 \mid b_\ell$, we must determine $\widetilde{\mathrm{Fr}}_\ell$ completely. Since $(\frac{\ell}{q}) = 1$, we have $(\frac{q}{\ell}) = 1$ and $(\frac{q^*}{\ell}) = 1$. Let $\alpha_\ell \in \mathbb{Z}$ such that $\alpha_\ell^2 \equiv q^* \bmod \ell$. From $\widetilde{\sigma}_q^{b_\ell}(\sqrt[4]{q^*}) = (-1)^{\frac{b_\ell}{2}}\sqrt[4]{q^*}$, we see $\widetilde{\mathrm{Fr}}_\ell = \widetilde{\sigma}_q^{b_\ell}$ if $(\frac{\alpha_\ell}{\ell}) = (-1)^{\frac{b_\ell}{2}}$, and $\widetilde{\mathrm{Fr}}_\ell = \widetilde{\sigma}_q^{b_\ell}\varepsilon$ if $(\frac{\alpha_\ell}{\ell}) = (-1)^{\frac{b_\ell}{2}+1}$. So when $\ell \equiv 1 \bmod 4$ and $\ell \equiv g^{b_\ell} \bmod q$ with $2 \mid b_\ell$, we have

$$L_\ell(\widetilde{\rho}_j, s) = \begin{cases} (1 - \zeta_{2(q-1)}^{jb_\ell}\ell^{-s})^{-2} & \text{if } (\frac{\alpha_\ell}{\ell}) = (-1)^{\frac{b_\ell}{2}} \\ (1 + \zeta_{2(q-1)}^{jb_\ell}\ell^{-s})^{-2} & \text{if } (\frac{\alpha_\ell}{\ell}) = (-1)^{\frac{b_\ell}{2}+1}. \end{cases}$$

Next we compute the local factors $L_2(\widetilde{\rho}_j, s)$ and $L_q(\widetilde{\rho}_j, s)$. When $(\frac{2}{q}) = -1$, we know from last section that 2 is ramified in $\widetilde{K}/K$. So $L_2(\widetilde{\rho}_j, s) = 1$

in this case. Now we assume $(\frac{2}{q}) = 1$. Since $I_2 =< \sigma_{-1} >$ and 2 is unramifed in $\widetilde{K}/K$, we have $\widetilde{I}_2 =< \widetilde{\sigma}_{-1} >$ or $\widetilde{I}_2 =< \widetilde{\sigma}_{-1}\varepsilon >$. The matrixes $I + \widetilde{\rho}_j(\widetilde{\sigma}_{-1})$ and $I + \widetilde{\rho}_j(\widetilde{\sigma}_{-1}\varepsilon)$ have rank 1. So $V^{\widetilde{I}_2}$ has dimension 1. Write $\mathrm{Fr}_2 = \sigma_2^{b_2}$ with $2 \mid b_2$. As in last section, we always assume $b_2 \equiv 2 \mathrm{mod} 4$ if $q \equiv 3 \mathrm{mod} 4$. Recall that $P_0$ be the set of the prime numbers of the form $A^2 + 64B^2$ with $A, B \in \mathbb{Z}$. Since $\widetilde{\rho}_j(\widetilde{\sigma}_2^{b_2}) = \zeta_{2(q-1)}^{jb_2} I$, by Prop.4.3 we have

$$L_2(\widetilde{\rho}_j, s) = \begin{cases} 1 - \zeta_{2(q-1)}^{jb_2} 2^{-s} & \text{if } \ q \notin P_0, 16 \nmid q^* - 1, \text{ or } q \in P_0, 16 \mid q^* - 1 \\ 1 + \zeta_{2(q-1)}^{jb_2} 2^{-s} & \text{if } \ q \in P_0, 16 \nmid q^* - 1, \text{ or } q \notin P_0, 16 \mid q^* - 1. \end{cases}$$

When $q \equiv 3 \mathrm{mod} 4$, we know that $q$ is ramified in $\widetilde{K}/K$. So $L_q(\widetilde{\rho}_j, s) = 1$ for odd $j$ in this case. Assume $q \equiv 1 \mathrm{mod} 4$. Since $I_q =< \sigma_q >$ and $q$ is unramifed in $\widetilde{K}/K$, we have $\widetilde{I}_q =< \widetilde{\sigma}_q >$ or $\widetilde{I}_2 =< \widetilde{\sigma}_q\varepsilon >$. Thus $V^{\widetilde{I}_q} = 0$ if $j \neq 0$, and $V^{\widetilde{I}_q}$ has dimension 1 if $j = 0$.

The Frobenious map $\mathrm{Fr}_q$ of $q$ in $G$ modulo $I_q$ is the identity map. So $\widetilde{\mathrm{Fr}}_q = 1$ or $\varepsilon$. In [Sect.5, 7] we have showed that $q$ splits in $\widetilde{K}/K$ if $q \equiv 1 \mathrm{mod} 8$ and is inertia if $q \equiv 5 \mathrm{mod} 8$. So $\widetilde{\mathrm{Fr}}_2 = 1$ if $q \equiv 1 \mathrm{mod} 8$ and $\widetilde{\mathrm{Fr}}_2 = \varepsilon$ if $q \equiv 5 \mathrm{mod} 8$. Thus we get

$$L_q(\widetilde{\rho}_j, s) = \begin{cases} 1 & \text{if } j \neq 0 \\ 1 - q^{-s} & \text{if } j = 0, \ q \equiv 1 \mathrm{mod} 8 \\ 1 + q^{-s} & \text{if } j = 0, \ q \equiv 5 \mathrm{mod} 8. \end{cases}$$

We have computed all the local factors. So we have

$$\begin{aligned} L(\widetilde{\rho}_j, s) =& (1 - u_q \zeta_{2(q-1)}^{jb_2} 2^{-s})^{-1} (1 - (-1)^{\frac{q-1}{4}} q^{-s})^{-n_j} \\ & \times \prod_{\ell \equiv 1, \ 2 \nmid b_\ell \text{ or } \ell \equiv 3, \ 2 \mid b_\ell} (1 - \zeta_{q-1}^{jb_\ell} \ell^{-2s})^{-1} \\ & \times \prod_{\ell \equiv 3, \ 2 \nmid b_\ell} (1 + \zeta_{q-1}^{jb_\ell} \ell^{-2s})^{-1} \prod_{\ell \equiv 1, \ 2 \mid b_\ell} (1 - u_\ell \zeta_{2(q-1)}^{jb_\ell} \ell^{-s})^{-2}, \end{aligned} \tag{5.2}$$

where $u_q = 1$ if $q \notin P_0, 16 \nmid q^* - 1$, or $q \in P_0, 16 \mid q^* - 1$ and $u_q = -1$ otherwise; $n_j = 0$ if $j \neq 0$ and $n_0 = 1$; and $u_\ell = (\frac{\alpha_\ell}{\ell})(-1)^{\frac{b_\ell}{2}}$. Here in the products, " $\equiv$ " means the congruence modulo 4.

**Theorem 5.1.** *Except for the Dirichlet L-functions, all Artin L-functions of the Galois extension* $\widetilde{K}/\mathbb{Q}$ *are explicitly given by (5.1) in the case* $q = 2$ *and by (5.2) in the case* $q$ *is odd, where in (5.2)* $0 \leq j < q-1, \ 2 \mid j$ *if* $q \equiv 1 \mathrm{mod} 4$ *and* $0 \leq j < q - 1, \ 2 \nmid j$ *if* $q \equiv 3 \mathrm{mod} 4$.

**5.3 A formula.** Let $\zeta_{\widetilde{K}}(s)$ and $\zeta_K(s)$ be the Dedekind zeta functions of $\widetilde{K}$ and $K$ respectively. By Artin's formula of the decomposition of Dedekind zeta functions, we have

$$\frac{\zeta_{\widetilde{K}}(s)}{\zeta_K(s)} = \prod_{\widetilde{\rho}_j} \prod_{\ell\text{: prime}} L_\ell(\widetilde{\rho}_j, s)^2,$$

where $\widetilde{\rho}_j$ runs over all 2-dimensional irreducible representations of $\widetilde{G}$. When $q = 2$, there is only one 2-dimensional irreducible representation of $\widetilde{G}$. So the square of (5.1) gives the formula. When $q$ is odd, by computing $\prod_{\widetilde{\rho}_j} L_\ell(\widetilde{\rho}_j, s)$, we get the following

**Corollary 5.2.** *For a prime number $\ell \neq q$, let $f_\ell = \frac{q-1}{\gcd(b_\ell, q-1)}$ be the order of $\ell \bmod q$ and let $g_\ell = \gcd(b_\ell, q-1) = \frac{q-1}{f_\ell}$. If $q \equiv 1 \bmod 4$, we have*

$$\begin{aligned}\frac{\zeta_{\widetilde{K}}(s)}{\zeta_K(s)} =&(1 - u_q^{f_2} 2^{-f_2 s})^{-g_2} (1 - (-1)^{\frac{q-1}{4}} q^{-s})^{-2} \prod_{\ell \equiv 1,\ 2 \nmid b_\ell \ or\ \ell \equiv 3} (1 - \ell^{-f_\ell s})^{-2g_\ell} \\ &\times \prod_{\ell \equiv 1,\ 2 \mid b_\ell} (1 - u_\ell^{f_\ell} \ell^{-f_\ell s})^{-2g_\ell},\end{aligned}$$

*and if $q \equiv 3 \bmod 4$, we have*

$$\begin{aligned}\frac{\zeta_{\widetilde{K}}(s)}{\zeta_K(s)} =&(1 + u_q^{f_2} 2^{-f_2 s})^{-g_2} \prod_{\ell \equiv 1,\ 2 \nmid b_\ell} (1 + \ell^{-f_\ell s})^{-2g_\ell} \prod_{\ell \equiv 3} (1 - \ell^{-2f_\ell s})^{-g_\ell} \\ &\times \prod_{\ell \equiv 1,\ 2 \mid b_\ell} (1 + u_\ell^{f_\ell} \ell^{-f_\ell s})^{-2g_\ell},\end{aligned}$$

*where $u_q$ and $u_\ell$ are as above.*

# References

[1] G. W. Anderson, *Kronecker-Weber plus epsilon*, Duke Math. J. **114**, 439-475 (2002).

[2] M. Daberkow, M. Pohst, *On integral bases in relative quadratic extensions*, Math. of Computation **65**, 319-329(1996).

[3] P. Das, *Algebraic Gamma monomials and double coverings of cyclotomic fields*, Trans. Amer. Math. Soc. **352**, 3557–3594 (2000).

[4] Y. Hu, L. Yin, *Arithmetic and representations of a non-abelian Galois number field*, Preprint, 2006.

[5] K. Ireland, M. Rosen, *A Classical Introduction to Modern Number Theory*, GTM 84, Springer-Verlag, 1982.

[6] J-P. Serre, *Linear Representations of finite groups*, GTM 42, Springer-Verlag, New York Inc. 1977.

[7] L. Yin and C. Zhang, *Arithmetic of quasi-cyclotomic fields*, Preprint, 2006.

[8] L. Yin and Q. Zhang, *All double coverings of cyclotomic fields*, Math. Zeit. **253**, 479-488(2006).